THERA Meeting *DESY*

19 October 2000 *Hamburg*

Prospects for Measurements of Diffractive Vector-Meson Photoproduction at High Momentum Transfer at THERA

- Recent Observations from ZEUS
- Open Questions for THERA

J. A. Crittenden

Physikalisches Institut
Universität Bonn

Diffractive Photoproduction of Vector Mesons

Elastic



Proton-dissociative



$\sqrt{s}$: Center–of–mass energy of e–p system
HERA (1996/97): 300 GeV

W: Center–of–mass energy of γ^*–p system
This analysis (ZEUS 1996/97 preliminary):
$80 < W < 120$ GeV

Q^2: Photon virtuality
This analysis: $Q^2 < 0.01$ GeV^2 ($\Rightarrow t \simeq -P_T^2(V)$)

t: Square of the momentum transfer
This analysis: $-11 < t < -1.5$ GeV^2
Proton-dissociative process dominates

M_N: Mass of the dissociated proton system
This analysis: $M_p^2 < M_N^2 < -100 * t$

Photoproduction Tagging at $W \simeq$ 100 GeV



$Q^2 < 0.01$ GeV^2

$23 < E_{e'} < 25$ GeV

$(\Rightarrow 80 < W < 120$ GeV$)$

ZEUS 1996/97 preliminary

Integrated luminosity: 24 pb^{-1}

ρ^0: $\simeq$ 18k

ϕ: $\simeq$ 2k

J/ψ: $\simeq$ 150

Candidate for ρ^0 Photoproduction at High $|t|$



$e^+p \to e^+N\,\rho^0 \to e^+N\,\pi^+\pi^-$ $\qquad$ $M_{\pi^+\pi^-} = 0.78$ GeV, $t = -10.0$ GeV^2, $W = 81$ GeV

Diffractive ρ^0 Photoproduction (I)

$$\gamma + p \rightarrow \rho^0 + N$$



Asymptotic region

Diffractive ρ^0 Photoproduction (II)



Scaling Property

Power dependence with

$n = 3$

matches the data fairly well

Diffractive ρ^0 Photoproduction (III)



Scaling with $(-t)^n$

Diffractive ϕ Photoproduction



Scaling with $(-t)^n$

Ratio of ϕ to ρ^0 Production



SU(3) flavor symmetry observed at high $|t|$

Diffractive J/ψ Photoproduction



Scaling with $(-t)^n$

Decay-Angle Analysis (I)



Choice of Z-axis:

"Helicity Frame"

Φ_h Angle between the electron scattering plane and the ρ^0 production plane

θ_h Polar angle of π^+ in the ρ^0 rest frame

ϕ_h Azimuthal angle of π^+

At low Q^2, Φ_h poorly measured, so integrate over it:

$$W(\cos\theta_h, \phi_h) = \frac{3}{4}\Big[\frac{1}{2}(1 - r^{04}_{00}) + \frac{1}{2}(3r^{04}_{00} - 1)\cos^2\theta_h - \sqrt{2}\,\mathrm{Re}(r^{04}_{10})\sin 2\theta_h \cos\phi_h - r^{04}_{1-1}\sin^2\theta_h \cos 2\phi_h \Big]$$

Decay-Angle Analysis (II)

$$r_{00}^{04} \simeq \frac{T_{01}^2}{T_{01}^2 + T_{11}^2 + T_{1-1}^2}$$

r_{00}^{04} measures the degree of longitudinal polarization



Tranverse polarization dominates

General Remarks

- These are the first measurements of vector-meson photoproduction at values of t comparable to the Q^2 values in deep inelastic scattering which led to the discovery of Bjorken scaling and charged proton constituents.

- This is presumably a strong interaction

- These values for t exceed the mass scales for ρ^0 and ϕ and Λ_{QCD} $\Rightarrow$ Asymptotic region
  The t dependence characterizes the interaction.

- We observe an extremely hard t spectrum
  $\Rightarrow$ low-order process

- Field theory for interaction without color exchange?

Conclusions I

High-t vector-meson photoproduction presents an opportunity to study the dynamics of diffractive processes

In the asymptotic region of momentum transfer

ρ^0 photoproduction exhibits a **very hard spectrum**

$$\frac{\mathrm{d}\sigma}{\mathrm{d}t} \propto (-t)^{-n} \quad n \simeq 3$$

The final-state ρ^0 are predominantly **transversely polarized**

The ratio of ϕ to ρ^0 exhibits **SU(3) flavor symmetry**

Conclusions II

At THERA the kinematics permit
reach to higher momentum transfer

The energy dependence is of
essential importance to clarify
the underlying physical process

Ratio of J/ψ to ρ^0 Production



Diffractive ϕ Photoproduction (I)

$$\gamma + p \rightarrow \phi + N$$



Asymptotic region

Diffractive ϕ Photoproduction (II)



Scaling Property

Power dependence with

$n = 3$

matches the data well

Diffractive J/ψ Photoproduction (I)

$$\gamma + p \rightarrow J/\psi + N$$



NOT asymptotic region

Diffractive J/ψ Photoproduction (II)



Scaling Property

Power dependence with

$n = 2$

matches the data fairly well