

The Evolution of the Singlet Structure Functions of the Proton and the Photon at Small x

J. Blümlein^a and A. Vogt^b

^a*DESY-Zeuthen
Platanenallee 6, D-15735 Zeuthen, Germany*

^b*Institut für Theoretische Physik, Universität Würzburg
Am Hubland, D-97074 Würzburg, Germany*

1. Basic Formalism
- (2. Non-Singlet Combinations)
3. $F_2^p(x, Q^2)$ and $F_L^p(x, Q^2)$
4. $F_2^\gamma(x, Q^2)$
5. Conclusions

1. Basic Formalism

$$q_{NS,i}^{\pm} = q_i \pm \bar{q}_i - \frac{1}{N_f} \sum_{r=1}^{N_f} (q_r \pm \bar{q}_r),$$

$$q_{NS}^{\text{val}} = \sum_{r=1}^{N_f} (q_r - \bar{q}_r),$$

$$q_S = \begin{pmatrix} \Sigma \\ g \end{pmatrix}, \quad \Sigma \equiv \sum_{r=1}^{N_f} (q_r + \bar{q}_r).$$

$$F_i(x, Q^2) = \sum_{r=1}^{2N_f} a_{ir} c_{i,r}(x, Q^2) \otimes q_r(x, Q^2) + a_{ig} c_{i,g}(x, Q^2) \otimes g(x, Q^2),$$

$$\frac{\partial q_{NS,i}^{\pm}(x, Q^2)}{\partial \ln Q^2} = P_{NS,i}^{\pm}(x, \alpha_s) \otimes q_{NS,i}^{\pm}(x, Q^2),$$

$$\frac{\partial q_S(x, Q^2)}{\partial \ln Q^2} = P_S(x, \alpha_s) \otimes q_S(x, Q^2).$$

$$\frac{d\alpha_s}{d \ln Q^2} = - \sum_{k=0}^{\infty} \alpha_s^{k+2} \beta_k.$$

$$P^{\pm}(x, \alpha_s) = \sum_{l=0}^{\infty} \alpha_s^{l+1} P_l^{\pm}(x),$$

$$P(x, \alpha_s) \equiv \begin{pmatrix} P_{qq}(x, \alpha_s) & P_{qg}(x, \alpha_s) \\ P_{gq}(x, \alpha_s) & P_{gg}(x, \alpha_s) \end{pmatrix} = \sum_{l=0}^{\infty} \alpha_s^{l+1} P_l(x),$$

$$c_{i,j}(x, Q^2) = \delta(1-x) \delta_{jq} + \sum_{l=1}^{\infty} \alpha_s^l c_{ij,l}(x).$$

$$\int_0^1 dx P_l^-(x) = 0,$$

$$\int_0^1 dx x \sum_i P_{ij,l}^{\text{unpol.}}(x) = 0,$$

$$\gamma_L(N, a_s) = -2 \begin{pmatrix} 0 & 0 \\ C_F/C_A & 1 \end{pmatrix} \gamma_L(N, \alpha_s).$$

$$\gamma(N, a_s) = -2 \int_0^1 dx x^{N-1} P(x, a_s).$$

$$\rho \equiv \frac{N}{\alpha_s} = 2\psi(1) - \psi(\gamma_L) - \psi(1 - \gamma_L) \equiv \chi[\gamma_L],$$

LIPATOV et al.
eq.

$$\gamma_L(N, a_s) = \sum_{k=1}^{\infty} g_k^{(0)} \left(\frac{\bar{\alpha}_s}{N} \right)^k.$$

$$\gamma_{\text{NL}}(N, \alpha_s) = -2 \begin{pmatrix} \frac{C_F}{C_A} [\gamma_{\text{NL}} - \frac{8}{3} a_s T_F] & \gamma_{\text{NL}} \\ \gamma_{gq, \text{NL}} & \gamma_{gg, \text{NL}} \end{pmatrix},$$

$$\begin{aligned} \gamma_{\text{NL}}^{\text{DIS}}(N, a_s) &= \gamma_{\text{NL}}^{Q_0}(N, a_s) R(\gamma_L) = T_F \frac{\alpha_s}{3\pi} \frac{2 + 3\gamma_L - 3\gamma_L^2}{3 - 2\gamma_L} \frac{[B(1 - \gamma_L, 1 + \gamma_L)]^3}{B(2 + 2\gamma_L, 2 - 2\gamma_L)} R(\gamma_L) \\ &= \mathcal{L} \frac{\alpha_s}{3\pi} T_F \sum_{k=1}^{\infty} g_k^{gq, (1)} \left(\frac{\bar{\alpha}_s}{N} \right)^k, \end{aligned}$$

CATANI 94
HAUTHMANN

with $B(x, y)$ denoting the Beta function and

$$R(\gamma) = \left[\frac{\Gamma(1 - \gamma)\chi(\gamma)}{\Gamma(1 + \gamma)\{-\gamma\chi'(\gamma)\}} \right]^{1/2} \exp \left[\gamma\psi(1) + \int_0^\gamma d\zeta \frac{\psi'(1) - \psi'(1 - \zeta)}{\chi(\zeta)} \right].$$

ALL THE ABOVE QUANTITIES ARE FUNCTIONS OF $\gamma_L(N)$.




Figure 1a: Real part of $\gamma_L(\rho)$ for $-3 \leq \text{Re} \rho \leq 3$ and $-5 \leq \text{Im} \rho \leq 5$. The labels of the curves $\gamma_L|_{\text{Re} \rho}$ refer to the order of their values for $|\text{Im} \rho| \sim 0$. The dash-dotted lines show the behavior of $\text{Re} \gamma_L$ at the singularities of the perturbative branch, $\text{Re} \rho = 4 \ln 2$ and -1.4105 . The branch points manifest by the 'roof' at $\gamma_L = 0.5$ and the corner points of $\gamma_L(\text{Re} \rho = -1.4105)$. The dashed lines mark a transition in γ_L for smaller values of $\text{Im} \rho$ at $\text{Re} \rho = 1.292$. The dotted lines illustrate the behavior of $\text{Re} \gamma_L$ outside the range of the singularities.




Figure 1b: Imaginary part of $\gamma_L(\rho)$ for $-3 \leq Re \rho \leq 3$ and $-5 \leq Im \rho \leq 5$. The notations are the same as in figure 1a.

N_f -dep. TERMS : γ_{gg} .

CAMICI
CIAFALONI 96

$$\gamma_{gg,NL}^{q\bar{q},Q_0} = -\alpha_s \frac{\chi_1^{(a)}(\gamma_L) + \chi_1^{(na)}(\gamma_L)}{\chi'(\gamma_L)} - \frac{C_F}{C_A} \gamma_{gg,NL}^{Q_0}.$$

$$\alpha_s \chi_1^{(a)}(\gamma_L) = \frac{2}{\gamma_L^2} \frac{C_F}{C_A} \gamma_{gg,NL}^{Q_0}(\gamma_L)$$

$$\alpha_s \chi_1^{(na)}(\gamma_L) = \frac{\alpha_s N_f}{6\pi} \left\{ \frac{1}{2} [\chi'(\gamma_L) + \chi^2(\gamma_L)] - \frac{5}{3} \chi(\gamma_L) \right\} - \frac{1}{\gamma_L^2} \frac{C_F}{C_A} \gamma_{gg,NL}^{Q_0}(\gamma_L).$$

$$\begin{aligned} \gamma_{gg,NL}^{q\bar{q},Q_0} = & -\frac{N_f \alpha_s}{6\pi} \left\{ 1 + \frac{23}{6} \frac{\bar{\alpha}_s}{N} + \left[\frac{71}{18} - \frac{\pi^2}{6} \right] \left(\frac{\bar{\alpha}_s}{N} \right)^2 + \left[\frac{233}{27} - \frac{13}{36} \pi^2 - 8\zeta(3) \frac{C_F}{C_A} \right] \left(\frac{\bar{\alpha}_s}{N} \right)^3 \right. \\ & + \left[\frac{1276}{81} - \frac{71}{108} \pi^2 + \frac{79}{3} \zeta(3) - \frac{7}{120} \pi^4 - \frac{52}{3} \zeta(3) \frac{C_F}{C_A} \right] \left(\frac{\bar{\alpha}_s}{N} \right)^4 \quad \uparrow ! \\ & + \left[\left(\frac{8384}{243} - \frac{233}{162} \pi^2 + \frac{284}{9} \zeta(3) - \frac{91}{720} \pi^4 + 2\zeta(5) - \frac{4}{3} \zeta(3) \pi^2 \right) \right. \\ & + \left. \left(\frac{4}{3} \zeta(3) \pi^2 - \frac{284}{9} \zeta(3) - 16\zeta(5) \right) \frac{C_F}{C_A} \right] \left(\frac{\bar{\alpha}_s}{N} \right)^5 \\ & + \left[\left(\frac{45928}{729} - \frac{638}{243} \pi^2 - \frac{65}{18} \zeta(3) \pi^2 - 2\zeta(3)^2 - \frac{497}{2160} \pi^4 + \frac{125}{3} \zeta(5) + \frac{2330}{27} \zeta(3) \right. \right. \\ & \left. \left. - \frac{31}{3024} \pi^6 \right) + \left(\frac{26}{9} \pi^2 \zeta(3) - \frac{104}{3} \zeta(5) - \frac{1864}{27} \zeta(3) - 80\zeta(3)^2 \right) \frac{C_F}{C_A} \right] \left(\frac{\bar{\alpha}_s}{N} \right)^6 \\ & \left. + O\left(\frac{\bar{\alpha}_s}{N} \right)^7 \right\}. \end{aligned}$$

In the DIS- \overline{MS} scheme $\gamma_{gg,NL}^{q\bar{q}}$ is given by, cf. [15],

$$\begin{aligned} \gamma_{gg,NL}^{q\bar{q},DIS} &= \gamma_{gg,NL}^{q\bar{q},Q_0} + \beta_0 \alpha_s^2 \frac{d \ln R(\alpha_s)}{d\alpha_s} + \frac{C_F}{C_A} [1 - R(\alpha_s)] \gamma_{gg,NL}^{Q_0} \\ &\equiv \alpha_s \sum_{k=1}^{\infty} \left[\frac{N_f}{6\pi} \left(\underline{d_{gg,k}^{q\bar{q},(1)}} + \frac{C_F}{C_A} \underline{d_{gg,k}^{q\bar{q},(2)}} \right) + \frac{\beta_0 \hat{r}_k}{4\pi} \right] \left(\frac{\bar{\alpha}_s}{N} \right)^{k-1}. \end{aligned}$$

\uparrow DIS.

F_L :

$$C_L^g = \frac{\alpha_s}{3\pi} T_F h_L(\gamma_L) R(\gamma_L) = \frac{2\alpha_s}{3\pi} T_F \sum_{k=1}^{\infty} c_k^L \left(\frac{\bar{\alpha}_s}{N} \right)^k,$$

$$C_L^q = \frac{C_F}{C_A} \left[C_L^g - \frac{2\alpha_s}{3\pi} T_F \right],$$

$$h_L(\gamma) = \left(\frac{1-\gamma}{3-2\gamma} \right) \frac{[B(1-\gamma, 1+\gamma)]^3}{B(2-2\gamma, 2+2\gamma)}.$$

k	$g_k^{(0)}$	$g_k^{qq,1}$	c_k^L
1	1.000000000E+0	1.000000000E+0	1.000000000E+0
2	0.000000000E+0	2.166666666E+0	-3.333333333E-1
3	0.000000000E+0	2.299510377E+0	2.132843711E+0
4	2.404113806E+1	8.271090891E+0	2.272314949E+0
5	0.000000000E+0	1.492492689E+1	0.434344619E+0
6	2.073855510E+1	2.922678142E+1	2.026429511E+1
7	1.733928958E+1	1.028116363E+2	2.303149346E+1
8	2.016698555E+0	1.948865992E+2	3.464487971E+1
9	3.988627732E+1	4.851003221E+2	2.650042403E+2
10	1.687465187E+2	1.524438954E+3	3.300383179E+2
11	6.998811177E+1	3.114512279E+3	8.503712548E+2
12	6.612526445E+2	8.583754389E+3	3.908491758E+3
13	1.945314152E+3	2.475708601E+4	5.674329386E+3
14	1.717675538E+3	5.474354877E+4	1.776800580E+4
15	1.064326485E+4	1.561954918E+5	6.219822925E+4
16	2.556678709E+4	4.269798325E+5	1.070279253E+5
17	3.678133240E+4	1.011107922E+6	3.514745865E+5
18	1.716848319E+5	2.893976929E+6	1.050584301E+6
19	3.753792748E+5	7.690417370E+6	2.103406452E+6
20	7.360245729E+5	1.919194565E+7	6.807468863E+6
k	$d_{gg,k}^{qq,(1)}$	$d_{gg,k}^{qq,(2)}$	\hat{r}_k
1	-1.000000000E+0	0.000000000E+0	0.000000000E+0
2	-3.833333333E+0	0.000000000E+0	0.000000000E+0
3	-2.299510376E+0	0.000000000E+0	0.000000000E+0
4	-5.065605818E+0	3.205485075E+0	9.616455224E+0
5	-3.523670351E+1	8.568702514E+0	-3.246969702E+0
6	-3.218245315E+1	1.835447655E+1	2.281241061E+1
7	-1.060268680E+2	8.632838009E+1	1.654162989E+2
8	-4.853159484E+2	1.924088636E+2	-2.469139930E+0
9	-5.806186371E+2	4.962344972E+2	7.458249428E+2
10	-2.176371931E+3	1.794742819E+3	2.784859262E+3
11	-7.553679737E+3	4.023320193E+3	1.505001272E+3
12	-1.158215080E+4	1.136559381E+4	1.818320928E+4
13	-4.328579102E+4	3.589638820E+4	4.899274185E+5
14	-1.269309428E+5	8.412529889E+4	6.109247725E+5
15	-2.392549581E+5	2.456097133E+5	3.984470167E+5
16	-8.469557573E+5	7.168572021E+6	9.205515787E+5
17	-2.262541206E+6	1.764587230E+6	1.783326920E+6
18	-4.974873276E+6	5.167844173E+6	8.347774614E+6
19	-1.648990863E+7	1.443009883E+7	1.842662795E+7
20	-4.222994214E+7	3.702246358E+7	4.535538189E+7

CAMICI, CIAPAL.



DES 97-106

JB, A.VOGT

16.6.

Table 1: The expansion coefficients in eqs. (29,34,38) and (40).

CAMICI + CIAFALONI:

5/97.

$$\alpha_s \chi_1^{gg} = \frac{C_A \alpha_s}{4\pi} \left[-\frac{11}{6} (\chi_0^2 + \chi_0') + \left(\frac{67}{9} - \frac{\pi^2}{3} \right) \chi_0 \right.$$

$$\left. + \left(6b_3 + \frac{\pi^2}{3\gamma(1-\gamma)} + \dots \right) \right]$$

$$- \left(\frac{\pi}{\sin \pi \gamma} \right)^2 \frac{\omega \pi \gamma}{3(1-2\gamma)} \left[11 + \frac{\gamma(1-\gamma)}{(1+2\gamma)(3-2\gamma)} \right]$$

1st STEPS:
K. ELUIS et al.

$$\begin{aligned}
 [U_1, R_0] &= R_1 + U_1, \\
 [U_2, R_0] &= R_2 + R_1 U_1 + 2U_2, \\
 &\vdots \\
 [U_k, R_0] &= R_k + \sum_{i=1}^{k-1} R_{k-i} U_i + k U_k \equiv \tilde{R}_k + k U_k.
 \end{aligned} \tag{52}$$

These equations can be solved recursively by applying the eigenvalue decomposition of the LO splitting function matrix, completely analogous to the NLO case of only U_1 in ref. [4]. One writes

$$R_0 = r_- e_- + r_+ e_+, \tag{53}$$

where r_- (r_+) stands for the smaller (larger) eigenvalue of R_0 ,

$$r_{\pm} = \frac{1}{2\beta_0} \left[P_{qq}^{(0)} + P_{gg}^{(0)} \pm \sqrt{(P_{qq}^{(0)} - P_{gg}^{(0)})^2 + 4P_{qg}^{(0)} P_{gq}^{(0)}} \right]. \tag{54}$$

The matrices e_{\pm} denote the corresponding projectors,

$$\begin{aligned}
 e_- &= \frac{1}{r_- - r_+} [R_0 - r_+ I], \\
 e_+ &= \frac{1}{r_+ - r_-} [R_0 - r_- I],
 \end{aligned} \tag{55}$$

with I the 2×2 unit matrix. Hence the LO evolution operator in eq. (50) can be written as

$$L(a, a_0, N) = e_-(N) \left(\frac{a_s}{a_0} \right)^{-r_-(N)} + e_+(N) \left(\frac{a_s}{a_0} \right)^{-r_+(N)}. \tag{56}$$

Inserting the obvious identity

$$U_k = e_- U_k e_- + e_- U_k e_+ + e_+ U_k e_- + e_+ U_k e_+ \tag{57}$$

into eq. (52), one finally obtains the expansion coefficients in eq. (51) as

$$U_k = -\frac{1}{k} [e_- \tilde{R}_k e_- + e_+ \tilde{R}_k e_+] + \frac{e_+ \tilde{R}_k e_-}{r_- - r_+ - k} + \frac{e_- \tilde{R}_k e_+}{r_+ - r_- - k}. \tag{58}$$

SOLUTION OF THE EVOLUTION eqs.

$$\begin{aligned} \frac{\partial q(a_s, N)}{\partial a} &= -\frac{1}{\beta_0 a_s} \left[P_0(N) + a_s \left(P_1(N) - \frac{\beta_1}{\beta_0} P_0(N) \right) \right. \\ &\quad \left. + a_s^2 \left(P_2(N) - \frac{\beta_1}{\beta_0} P_1(N) + \left\{ \left(\frac{\beta_1}{\beta_0} \right)^2 - \frac{\beta_2}{\beta_0} \right\} P_0(N) \right) + \dots \right] q(a_s, N) \\ &\equiv -\frac{1}{a_s} \left[R_0(N) + \sum_{k=1}^{\infty} a_s^k R_k(N) \right] q(a_s, N). \end{aligned}$$

$$R_0 = \frac{1}{\beta_0} P_0,$$

$$R_k = \frac{1}{\beta_0} P_k - \sum_{i=1}^k \frac{\beta_i}{\beta_0} R_{k-i}$$

$$[R_{k \geq 1}, R_0] \neq 0$$

$$q^{LO}(a_s, N) = \left(\frac{a_s}{a_0} \right)^{-R_0(N)} q(a_0, N) \equiv L(a_s, a_0, N) q(a_0, N),$$

$$\begin{aligned} q(a_s, N) &= U(a_s, N) L(a_s, a_0, N) U^{-1}(a_0, N) q(a_0, N) \\ &= \left[1 + \sum_{k=1}^{\infty} a_s^k U_k(N) \right] L(a_s, a_0, N) \left[1 + \sum_{k=1}^{\infty} a_0^k U_k(N) \right]^{-1} q(a_0, N). \end{aligned}$$

SUBLEADING TERMS:

$$\gamma_{99,1}^{Dis} = \frac{N_f = 4}{N} - \frac{123.3}{N} + 405.9$$

$$\gamma_{98,1}^{Dis} = - \frac{277.3}{N} + 846.2$$

$$\gamma_{89,1}^{Dis} = \frac{91.26}{N} - 453.5$$

$$\gamma_{88,1}^{Dis} = \frac{245.3}{N} - 988.3$$

MODEL:

A: $\gamma_{ij} \rightarrow \gamma_{ij} - \delta_{ij} \gamma_{ij} (1, a_s)$

B: $\gamma_{ij} \rightarrow \gamma_{ij} \cdot (1 - N)$

C: $\gamma_{ij} \rightarrow \gamma_{ij} (1 - N)^2$


D: $\gamma_{ij} \rightarrow \gamma_{ij} (1 - 2N + N^2)$.

AN EXAMPLE IN NLO:


$$\delta F_3^{NLO} = \frac{\alpha_s \cdot C_3 \otimes x(u_V + d_V)}{x(u_V + d_V)}$$

JB, A. VOGT

CF. ALSO
KODAIRA et al
'97



— full
- - - Lx_1
... Lx_2
- · - Lx_3
· · · Lx_4 !





$$Lx_1 = Lx$$

$$Lx_2 = Lx + NLx$$

$$Lx_3 = Lx + NLx + NNLx$$





⋮



3. $F_2^P(x, Q^2)$, $F_L^P(x, Q^2)$



.... $C_L^{NLO} + C_L^{>A} \cdot (1-2N)$


F_L IS DRIVEN BY THE COEFFICIENT FUNCTION (NOT BY THE PDF'S).






$$\gamma_0(1 - 2N + N^2)$$

JB, A. VOGT



NO
ESSENTIAL
MODIFICATION.



- NLX resummation
- .- NLX · (1-N)
- NLO

4. $F_2^\gamma(x, Q^2)$

$$\begin{pmatrix} \dot{q}^\gamma \\ \dot{\bar{q}}^\gamma \\ \dot{G}^\gamma \\ \dot{\Gamma}^\gamma \end{pmatrix} = \frac{\alpha}{2\pi} \begin{pmatrix} P_{qq} \\ P_{q\bar{q}} \\ P_{Gq} \\ P_{\gamma q} \end{pmatrix} \otimes \Gamma^\gamma + \begin{cases} \frac{\alpha_s}{2\pi} P_{ij} \otimes P_j \\ \frac{\alpha}{2\pi} P_{\gamma j} \otimes P_j \end{cases} \quad i, j = 1, 3.$$

$$\Gamma^{\gamma,0} = \delta(1-x) + O(\alpha). \quad q^\gamma = \bar{q}^\gamma$$

SOLVE :

$$\dot{q}^\gamma = \frac{\alpha}{2\pi} K_q + \frac{\alpha_s}{2\pi} \left[\sum_{k=1}^{N_f} (P_{qq} + P_{q\bar{q}}) \otimes q^\gamma + P_{qg} \otimes G^\gamma \right]$$

$$\dot{G}^\gamma = \frac{\alpha}{2\pi} K_g + \frac{\alpha_s}{2\pi} \left[\sum_{k=1}^{N_f} (P_{gq} + P_{g\bar{q}}) \otimes q^\gamma + P_{gg} \otimes G^\gamma \right]$$

INHOM. EVOLUTION EQUS.

$$K_i = \sum_{k=0}^{\infty} \left(\frac{\alpha_s}{2\pi} \right)^k K_{i,k}$$

$$P_{ij} = \sum_{k=0}^{\infty} \left(\frac{\alpha_s}{2\pi} \right)^k P_{ij}^{(k)}$$

→ EXPRESS F_2^γ & EXPAND.

• SCHEME INDEP. REQUESTS :

$$K_0 = \text{inv.}$$

$$K_0 C_1 + K_1 - \delta \cdot P_0 C_{\gamma 1} = \text{inv.}$$

$$K_2 + C_2 K_0 + C_1 K_1 - \delta \cdot (P_0 C_{\gamma 2} + P_1 C_{\gamma 1} + [C_1, P_0] C_{\gamma 1})$$

$$+ \delta \beta_0 C_1 C_{\gamma 1} - \delta \beta_0 C_{\gamma 2} = \text{inv.}$$

etc.

Dis γ : NO NLX terms in K_n ($C_i \sim \frac{1}{(N-1)^{i-1}}$)

$K_{\text{Dis}, \gamma, \text{in}} = 0 \mid \text{NLX}$. TREAT IN NLO only.

$$C_{\gamma i} \sim \frac{1}{(N-1)^{i-1}}$$

PRODUCTS $k_i C_j, C_i C_j$:
 \rightarrow NO CONTRIB. IN NLX.

NLX:

$$\text{inv} = k^n - \delta \left(\underbrace{P_0 C_{\gamma, n} + \dots}_{\text{not NLX}} P_{n-1} C_{\gamma, 1} \right) - \delta \beta_0 \theta \cdot (N-2) \cdot C_{n, \gamma}$$

$x \rightarrow 0$

$C_{\gamma, n}$ ONLY FOR $N=1$, NLX

$$\text{inv} \Big|_{x \rightarrow 0} = k_n - \delta \cdot P_{n-1}^{\text{q}\gamma} C_{\gamma, 1}$$

\uparrow NO HADRONIC
 IMPACT, IF $\overline{\text{MS}} \rightarrow \text{DIS}$

$$R_{\overline{\text{MS}}, n} \Big|_{x \rightarrow 0} = \frac{1}{2} P_{\text{q}\gamma, 0} (N=1) \cdot P_{\text{g}\gamma, n-1} \Big|_{x \rightarrow 0}$$



\leftarrow Constant!

$$\rightarrow R_{\text{DIS}, n} \Big|_{x \rightarrow 0} = 0.$$

I.E. ONLY THE HADRONIC CONTR. ARE LEFT
 IN THIS WIDELY USED SCHEME.

PHOTON STRUCTURE FUNCTIONS

JB, AVOGT



GRV₈; DIS

- HADRONIC PART DOMINATES
- UP TO NLX THE INHOMOGENEITY DOES RECEIVE RESUMMED CONTRIBUTIONS IN THE DIS-X SCHEME.

5. Conclusions

- AN UPDATE OF THE SMALL x RESUMMATION FOR F_2 and F_L WAS PERFORMED.
- SUBLEADING TERMS IN $\left(\frac{\alpha_s}{N}\right)^k (\alpha_s N)^a$ ARE AS IMPORTANT. \rightarrow NEED TO BE CALCULATED.
 \rightarrow ANOMALOUS DIMENSIONS & COEFFICIENT FCTS.
- DIS SCHEME: F_L DEPENDS ON C_L^g MOSTLY,
 g - 'RATHER' STABLE!

$$\delta C_L^g \leftrightarrow \delta F_L!$$

- THE TERMS $\propto N_f$ IN γ_{gg} WERE INCLUDED. THEIR NUMERICAL EFFECT IS SMALL ($< 1\%$). STILL TO WAIT UNLESS THE COMPLETE NLO RESUMMED γ_{gg} & γ_{gq} BECOME AVAILABLE.
- THE SMALL x CONTRIBUTIONS TO F_2^{γ} WERE CALCULATED.

THE POINTLIKE TERM DOES NOT RECEIVE RESUMMED CORRECTIONS IN NLX IN THE DIS SCHEME.

THE CORRECTIONS ARE DUE TO THE HADRONIC PARTS.




Figure 1a: Real part of the perturbative branch of γ_L as a function of the real and imaginary part of $\rho = N/\bar{\alpha}_s$. The full lines correspond to $Re \rho = 1, 0, -1$ and the dash-dotted lines are the contours through the singularities.




Figure 1b: Imaginary part of $\gamma_L(\rho)$ for $-3 \leq \text{Re } \rho \leq 3$ and $-5 \leq \text{Im } \rho \leq 5$. The notations are the same as in figure 1a.

